
GCSE MATHEMATICS

Aiming for Grade 7

REVISION BOOKLET

Exam Dates:



Contents

	Page:
Number:	_
Surds	3
Bounds calculations	7
Algebra:	
Parallel and perpendicular graphs	11
Transformations of graphs	16
Algebraic fractions - simplifying	20
Algebraic fractions – solving	23
Solving quadratic inequalities	26
Shape, Space and Measure:	
Circle theorems	28
Vectors	32
Sine and cosine rules	40
Data Handling:	
Cumulative frequency and box plots	44
Histograms	51
Probability:	
Set theory	59
Ratio and Proportion:	
Proportion	62
Percentages – compound interest	66
Percentages – reverse	68

Surds

Things to remember:

- √ means square root;
- To simplify surds, find all its factors;
- To rationalise the denominator, find an equivalent fraction where the denominator is rational.

Questions:

1. Work out

$$\frac{(5+\sqrt{3})(5-\sqrt{3})}{\sqrt{22}}$$

Give your answer in its simplest form.

											(1	Ī	()	t	ć	1	I	,	3	r	r	1	a	ľ	·	<	S	;)

2. (a) Rationalise the denominator of $\frac{1}{\sqrt{3}}$

	(1)

(b) Expand $(2 + \sqrt{3})(1 + \sqrt{3})$ Give your answer in the form $a + b\sqrt{3}$ where a and b are integers.



(Total 3 marks)

3. (a) Rationalise the denominator of $\frac{1}{\sqrt{7}}$

(2)

(b) (i) Expand and simplify $(\sqrt{3} + \sqrt{15})^2$ Give your answer in the form $a + b\sqrt{3}$ where a and b are integers.

.....

(ii) All measurements on the triangle are in centimetres.ABC is a right-angled triangle.k is a positive integer.




Diagram **NOT** accurately drawn


Find the value of *k*.

k =

(Total 7 marks)

4.	Expa	and and simplify $(\sqrt{3} - \sqrt{2})(\sqrt{3} - \sqrt{2})$	
		1/	(Total 2 marks)
5.	(a)	Write down the value of $49^{1/2}$	(1)
	(b)	Write $\sqrt{45}$ in the form $k\sqrt{5}$, where k is an integer.	
			(1) (Total 2 marks)
6.	Write	$\frac{\sqrt{18+10}}{\sqrt{2}}$ in the form $a+b\sqrt{3}$ where a and b are in	

7. Expand and simplify $(2 + \sqrt{3})(7 - \sqrt{3})$ Give your answer in the form $a + b\sqrt{3}$ where a and b are integers.



8. Rationalise the denominator of $\frac{(4+\sqrt{2})(4-\sqrt{2})}{\sqrt{7}}$ Give your answer in its simplest form.



9. Show that $\frac{(4-\sqrt{3})(4+\sqrt{3})}{\sqrt{13}}$ simplifies to $\sqrt{13}$

(Total for question = 2 marks)

Bounds Calculations

Things to remember:

- Calculating bounds is the opposite of rounding they are the limits at which you would round up instead of down, and vice versa.
- When dividing bounds, UB = UB ÷ LB and LB = LB ÷ UB

Give your answer correct to 3 decimal places.

You must show your working.

Oi	ıes	ti	<u>_</u>	n	c	•
w	163	LI	v		J	

Ques	งแบบร	•	
1.	A pie (a)	ece of wood has a length of 65 centimetres to the n What is the least possible length of the piece of v	
	(b)	What is the greatest possible length of the piece	of wood?
			(1) (Total for Question is 2 marks)
2.	Chel (a)	lsea's height is 168 cm to the nearest cm. What is Chelsea's minimum possible height?	
	(b)	What is Chelsea's maximum possible height?	cm (1)
			cm (1)
		V.	(Total for Question is 2 marks)
3.		$\frac{V}{R}$ 250 correct to the nearest 5 3900 correct to the nearest 100	
	Worl	k out the lower bound for the value of /	

(Total for question = 3 marks)

4. Here is a solid bar made of metal.

The bar is in the shape of a cuboid.

The height of the bar is *h* cm.

The base of the bar is a square of side d cm.

The mass of the bar is $M \log M$

d = 8.3 correct to 1 decimal place.


M = 13.91 correct to 2 decimal places.

h = 84 correct to the nearest whole number.

Find the value of the density of the metal to an appropriate degree of accuracy.

Give your answer in g/cm³.

You must explain why your answer is to an appropriate degree of accuracy.



(Total for question = 5 marks)

5. Steve travelled from Ashton to Barnfield.

He travelled 235 miles, correct to the nearest 5 miles.

The journey took him 200 minutes, correct to the nearest 5 minutes.

Calculate the lower bound for the average speed of the journey.

Give your answer in **miles per hour**, correct to 3 significant figures.

You must show all your working.

mŗ	
(Total for question = 4 mark	S)

6.	The value of p is 4.3
	The value of q is 0.4

Both p and q are given correct to the nearest 0.1

(a) Write down the lower bound for p.

(1)

$$r = p + \frac{1}{q}$$

7.

(b) Work out the upper bound for *r*. You must show all your working.


(3)

(Total for question = 4 marks)

$$m = \frac{\sqrt{s}}{t}$$
 $s = 3.47$ correct to 3 significant figures $t = 8.132$ correct to 4 significant figures

By considering bounds, work out the value of \emph{m} to a suitable degree of accuracy. Give a reason for your answer.

8. *a* is 8.3 cm correct to the nearest mm *b* is 6.1 cm correct to the nearest mm



Calculate the upper bound for *c*. You must show your working.

	cm
(Total for question = 4 mark	เร


Parallel and Perpendicular Graphs

Things to remember:

- The general equation of a linear graph is given by y = mx + c, where m is the gradient and c is the y-intercept.
- Parallel graphs have the same gradient.
- Gradients of perpendicular graphs have a product of -1.

Questions:

1. The diagram shows a straight line, L₁, drawn on a grid.




A straight line, L_2 , is parallel to the straight line L_1 and passes through the point (0, -5). Find an equation of the straight line L_2 .

(Total for Question is 3 marks)

2. The straight line **L** has equation y = 2x - 5Find an equation of the straight line perpendicular to **L** which passes through (-2, 3).

(Total for Question is 3 marks)

3. In the diagram, *ABC* is the line with equation $y = -\frac{1}{2}x + 5$ AB = BC*D* is the point with coordinates (-13, 0)




Find an equation of the line through A and D.


(Total for question = 5 marks)

4. Here are the graphs of 6 straight lines.


Graph A




Graph B




Graph C




Graph ${\bf D}$



Graph E



Graph F




Match each of the graphs A, B, C, D, E and F to the equations in the table.

Equation	$y = \frac{1}{2}x + 2$	y = 2x - 2	$y = -\frac{1}{2}x + 2$	y = -2x - 2	y = 2x + 2	$y = -\frac{1}{2}x - 2$
Graph						

(Total for Question is 3 marks)

5. In the diagram, A is the point (-2, 0) B is the point (0, 4) C is the point (5, -1)



Find an equation of the line that passes through C and is perpendicular to AB.

(Total for Question is 4 marks)

6. Find an equation of the straight line that is perpendicular to the straight line x + 2y = 5 and that passes through the point (3, 7).

(Total for Question is 4 marks)

*7.	A and B are straight lines. Line A has equation $2y = 3x + 8$ Line B goes through the points $(-1, 2)$ and $(2, 8)$	
	Do lines A and B intersect? You must show all your working.	
		(Total for Question is 3 marks)
8.	A straight line, L , is perpendicular to the line with equation. The point with coordinates (6, 3) is on the line L . Find an equation of the line L .	y = 1 - 3x.
	•••••	(Total for Question is 3 marks)

Transformations of graphs


Things to remember:

- f(x) means the function of x.
- -f(x) is a reflection in the x-axis.
- f(-x) is a reflection in the y-axis.
- f(x a) is a translation in the x-axis, a units.
- f(x) + b is a translation in the y-axis, b units.
- cf(x) is an enlargement in the y-axis, scale factor c.
- f(dx) is an enlargement in the x-axis, scale factor $\frac{1}{d}$.

Questions:


1. y = f(x)

The graph of y = f(x) is shown on the grid.



(a) On the grid above, sketch the graph of y = -f(x).

The graph of y = f(x) is shown on the grid.



The graph **G** is a translation of the graph of y = f(x).


(b) Write down the equation of graph **G**.

(2) (Total for Question is 3 marks)


(2)

16

2. The graph of y = f(x) is shown on both grids below.




(a) On the grid above, sketch the graph of y = f(-x)




(b) On this grid, sketch the graph of y = -f(x) + 3

(1) (Total for question = 2 marks)

- The graph of y = f(x) is shown on each of the grids. (a) On this grid, sketch the graph of y = f(x 3)3.




(b) On this grid, sketch the graph of y = 2f(x)



(Total for Question is 4 marks)

(2)


The graph of y = f(x) is shown on the grid. 4.



On the grid above, sketch the graph of y = f(x + 3)(a)

(2)

The graph of y = g(x) is shown below.



The graph **G** is the reflection of y = g(x) in the *x*-axis. (b) Write down an equation of graph **G**.

(1)

(Total for question = 3 marks)

Algebraic Fractions - Simplifying

Things to remember:

- Factorise the numerator and denominator;
- Cancel common factors;
- Then add/subtract/multiply divide if necessary.

Questions:

1. Simplify
$$\frac{p^2-9}{2p+6}$$

(Total 3 marks)

2. Simplify fully $\frac{6x^2 + 3x}{4x^2 - 1}$

(Total 3 marks)

3. Simplify $\frac{x^2 + 2x + 1}{x^2 + 3x + 2}$

(Total 3 marks)

4. Simplify fully
$$\frac{x^2 + x - 6}{x^2 - 7x + 10}$$

5. Simplify fully
$$\frac{x^2 - 8x + 15}{2x^2 - 7x - 15}$$

6. Simplify fully
$$\frac{2x^2+3x+1}{x^2-3x-4}$$

7. (a) Simplify
$$\frac{2x+4}{x^2+4x+4}$$

(b) Write $\frac{1}{x+4} + \frac{2}{x-4}$ as a single fraction in its simplest form.

(3) (Total 6 marks)

8. Simplify fully
$$\frac{x+3}{4} + \frac{x-5}{3}$$

Algebraic fractions - solving

Things to remember:

- Multiply every term by the product of the denominators;
- Solve to find x.

Questions:

1. Solve
$$\frac{5(2x+1)}{3} = 4x + 7$$

X = (Total 3 marks)

 $\frac{5}{x+2} = \frac{4-3x}{x-1}$ Show that the equation 2. (a)

can be rearranged to give $3x^2 + 7x - 13 = 0$

Solve $3x^2 + 7x - 13 = 0$ (b) Give your solutions correct to 2 decimal places. (3)

x = or *x* =

(Total 6 marks)

3. Solve the equation
$$\frac{x}{2x-3} + \frac{4}{x+1} = 1$$

4. Solve the equation
$$\frac{3}{x+3} - \frac{4}{x-3} = \frac{5x}{x^2-9}$$

5. (a) Solve
$$\frac{3}{x} + \frac{3}{2x} = 2$$

(b) Using your answer to part (a), or otherwise, solve
$$\frac{3}{(y-1)^2} + \frac{3}{2(y-1)^2} = 2$$

(3) (Total 5 marks)

Solving Quadratic Inequalities

Things to remember:

Start by solving the quadratic to find the values of x, then sketch the graph to determine the inequality.

Questions:

Solve

 $x^2 > 3x + 4$

(Total for question = 3 marks)

Solve the inequality $x^2 > 3(x+6)$ 2.

$$x^2 > 3(x+6)$$

(Total for question = 4 marks)

Solve the inequality 3.


$$x^2 + 5x > 6$$

(Total for question = 3 marks)


4.	Solve the inequality	x² - 2x - 8 < 0	
			(Total for question = 3 marks)
5.	Solve the inequality	x² - x ≥ 12	
			(Total for question = 3 marks)
6.	Solve the inequality	$x^2 \le 4(2x + 5)$	
			(Total for question = 4 marks)

Circle theorems


Things to remember:




The angle at the centre is twice the angle at the circumference.




The angle in a semicircle is 90°.




Angles subtended by the same arc are equal.




Opposite angles in a cyclic quadrilateral sum to 180°.



Tangents from a point are equal.



A tangent is perpendicular to a radius.



Angles in alternate segments are equal.

Questions:

1.




Diagram **NOT** accurately drawn

P is a point on the circumference of the circle, centre O.

- PQ is a tangent to the circle.
- Write down the size of angle OPQ.

																													(
• • • • •	• • •	• • •	• • •	• •	• •	• •	• •	• •	•	• •	• •	• •	• •	• •	•	• •	• •	•	• •	• •	•	• •	• •	•	• •	• •	• •	• • • •	

(ii)	Give a reason for your answer.

(Total 2 marks)




Diagram NOT accurately drawn

Α,	B and C	Care	points	on t	he	circum	ference	of a	circle,	centre	0

AC is a diameter of the circle.

(a) (i) Write down the size of angle *ABC*.

(ii) Give a reason for your answer.




Diagram **NOT** accurately drawn

D, E and F are points on the circumference of a circle, centre O. Angle $DOF = 130^{\circ}$.

- (b) (i) Work out the size of angle *DEF*.
 - (ii) Give a reason for your answer.

(2) (Total 4 marks)

(2)

3.




Diagram **NOT** accurately drawn

A and B are points on the circumference of a circle, centre O.

PA and PB are tangents to the circle.

Angle APB is 86°.

Work out the size of the angle marked x.

(Total 2 marks)

4.




Diagram **NOT** accurately drawn

In the diagram, A, B, C and D are points on the circumference of a circle, centre O. Angle $BAD = 70^{\circ}$.

Angle $BOD = x^{\circ}$.

Angle $BCD = y^{\circ}$.

(a)	(1)	vvork out the value of x.	

	(2)
()		
(11)	Give a reason for your answer.	


Work out the value of y. (b) (i)

(ii)	Give a reason for your answer.	o

(Total 4 marks)

(2)

5.



DCO is a straight line.

DA is a tangent to the circle. Angle $ADO = 36^{\circ}$

(a) Work out the size of angle AOD.	
-------------------------------------	--

			(3) (Total 5 marks)
	(ii)	Give a reason for your answer.	
(b)	(i)	Work out the size of angle ABC.	c
			(2)
(α)	*****	it out the size of angle 710b.	c

Vectors

Things to remember:

- Use the letter provided in the question.
- Going against the arrow is a negative.
- Vectors need to be written in bold or underlined.
- They can be manipulated similarly to algebra.

Questions:

The diagram shows a regular hexagon ABCDEF with centre O. 1.




Diagram NOT accurately drawn

 $\overrightarrow{OA} = 6a$ $\overrightarrow{OB} = 6\mathbf{b}$

Express in terms of a and/or b (a)

AB(i)

 \overrightarrow{EF} (ii)

(2)

X is the midpoint of BC.

Express \overline{EX} in terms of **a** and/or **b** (b)


(2)

Y is the point on AB extended, such that AB : BY = 3:2

Prove that *E*, *X* and *Y* lie on the same straight line.

(Total 7 marks)

2. T is the point on PQ for which PT: TQ = 2:1.



OPQ is a triangle.

$$\overrightarrow{OP} = \mathbf{a} \text{ and } \overrightarrow{OQ} = \mathbf{b}.$$

(a) Write down, in terms of **a** and **b**, an expression for \overrightarrow{PQ} .

\overrightarrow{PQ}	=	 	 	 		 	 	 			 						 		 	 		
																				(1	1

(b) Express \overrightarrow{OT} in terms of \mathbf{a} and \mathbf{b} . Give your answer in its simplest form.

$$\overrightarrow{OT}$$
 =(2)
(Total 3 marks)

3. *OABC* is a parallelogram.




Diagram **NOT** accurately drawn

P is the point on *AC* such that $AP = \frac{2}{3}AC$.

$$\overrightarrow{OA} = 6a. \overrightarrow{OC} = 6c.$$

(a) Find the vector \overrightarrow{OP} . Give your answer in terms of **a** and **c**.

	(3)

The midpoint of CB is M.

(b) Prove that *OPM* is a straight line.

(2) (Total 5 marks) 4. OPQ is a triangle.

R is the midpoint of OP.

S is the midpoint of PQ. $\overrightarrow{OP} = \mathbf{p}$ and $\overrightarrow{OQ} = \mathbf{q}$




Diagram **NOT** accurately drawn

(i) Find \overrightarrow{OS} in terms of **p** and **q**.

\overrightarrow{OS}			
CD	=	 	

(ii) Show that RS is parallel to OQ.

(Total 5 marks)


OPQR is a trapezium with PQ parallel to OR. 5.

$$\overrightarrow{OP} = 2\mathbf{b}$$

$$\overrightarrow{PQ}$$
 = 2a \overrightarrow{OR} = 6a

$$\overrightarrow{OR} = 6a$$

M is the midpoint of PQ and N is the midpoint of OR.



Find the vector \overrightarrow{MN} in terms of **a** and **b**. (a)



X is the midpoint of MN and Y is the midpoint of QR.

Prove that XY is parallel to OR. (b)

(Total 4 marks)

ABCD is a straight line. 6.




Diagram **NOT** accurately drawn

O is a point so that $\overrightarrow{OA} = \mathbf{a}$ and $\overrightarrow{OB} = \mathbf{b}$.

B is the midpoint of AC.


C is the midpoint of AD.

Express, in terms of \mathbf{a} and \mathbf{b} , the vectors

 \overrightarrow{AC} (i)

(ii)	OD
()	

7. Diagram **NOT** accurately drawn



ABCD is a parallelogram. AB is parallel to DC.

AD is parallel to BC.

$$\stackrel{
ightarrow}{AB}={\bf p}$$

$$\stackrel{
ightarrow}{AD}={\bf q}$$


Express, in terms of p and q (a)

- $\stackrel{\rightarrow}{AC}$ (i)
- $\stackrel{
 ightarrow}{BD}$ (ii)



(2)

Diagram **NOT** accurately drawn



AC and BD are diagonals of parallelogram ABCD. AC and BD intersect at T.

Express \overrightarrow{AT} in terms of **p** and **q**. (b)

(Total 3 marks)


8. Diagram **NOT** accurately drawn

OAB is a triangle.

B is the midpoint of *OR*.

Q is the midpoint of AB.

$$\overrightarrow{OP}$$
 = 2a \overrightarrow{PA} = a \overrightarrow{OB} = b



- (a) Find, in terms of **a** and **b**, the vectors
 - (i) \overrightarrow{AB} ,
 - (ii) \overrightarrow{PR} ,

		\overrightarrow{PQ} .	(iii)	
(4	 			<i>(</i> 1. \

(b) Hence explain why *PQR* is a straight line.

	(2)
The length of PQ is 3 cm.	•

(c) Find the length of *PR*.

......cm

(Total 7 marks)


Sine and Cosine Rules

Things to remember:

• For any triangle ABC, $a^2 = b^2 + c^2 - 2bc \cos A$

• For any triangle ABC, $\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$

For any triangle ABC, area = ½ a b sinC



Questions:

1. Diagram **NOT** accurately drawn

ABC is a triangle.

D is a point on AC.

Angle $BAD = 45^{\circ}$

Angle $ADB = 80^{\circ}$


AB = 7.4 cm

DC = 5.8 cm

Work out the length of BC.

Give your answer correct to 3

significant figures.



.....cm (Total for question = 5 marks)

2. Diagram **NOT** accurately drawn

ABC is a triangle.

AB = 8.7 cm.


Angle $ABC = 49^{\circ}$.

Angle $ACB = 64^{\circ}$.

Calculate the area of triangle ABC.


Give your answer correct to 3 significant

figures.




......cm²
(Total for Question is 5 marks)

ABCD is a quadrilateral.
 Diagram NOT accurately drawn
 Work out the length of DC.
 Give your answer correct to 3 significant figures.




 				cn
(Total	for Qu	uestio	n is 6	marks

Diagram NOT accurately drawn
 ABC is an isosceles triangle.
 Work out the area of the triangle.
 Give your answer correct to 3 significant figures.




......cm²
(Total for Question is 4 marks)

5. Diagram NOT accurately drawn The diagram shows triangle LMN. Calculate the length of LN. Give your answer correct to 3 significant figures.



VABCD is a solid pyramid.
 ABCD is a square of side 20 cm.
 The angle between any sloping edge and the plane ABCD is 55°
 Calculate the surface area of the pyramid.
 Give your answer correct to 2 significant figures.




(Total for Question is 5 marks)

.....cm²
(Total for question = 5 marks)

7. The diagram shows triangle ABC. The area of triangle ABC is $k\sqrt{3}$ cm².

Find the exact value of *k*.



k =			
	Total for c	question =	7 marks


8. Diagram **NOT** accurately drawn

AC = 9.2 mBC = 14.6 m

Angle $ACB = 64^{\circ}$

(a) Calculate the area of the triangle *ABC*.
Give your answer correct to 3

significant figures.



..... m²

(b) Calculate the length of *AB*.

Give your answer correct to 3 significant figures.

.....(3)

(Total for Question is 5 marks)

Cumulative frequency and box plots

Things to remember:

- Use a running total adding on to complete the cumulative frequency column;
- Plot at the end of the group;
- Join up with a smooth curve;
- To find the median find the value half way down the cumulative frequency, draw across to the line and then vertically down to find the value – always show these working lines;
- To find the interquartile range find the upper quartile and the lower quartile and subtract them.

 Walking Market Control of the Con
- To draw a box plot -
- When comparing box plots, use the median and the IQR and keep words consistent with the question.




Questions:

1. The table shows information about the heights of 40 bushes.

Height (h cm)	Frequency	Cumulative Frequency
170 ≤ <i>h</i> < 175	5	
175 ≤ <i>h</i> < 180	18	
180 ≤ <i>h</i> < 185	12	
185 ≤ <i>h</i> < 190	4	
190 ≤ <i>h</i> < 195	1	

(a) Complete the cumulative frequency table above.

(b) On the grid, draw a cumulative frequency graph for your table.



(2) (Total 3 marks)

(1)

44

2. The table gives information about the ages of 160 employees of an IT company.

Age (A) in years	Frequency	Cumulative Frequency
15 < <i>A</i> ≤ 25	44	
25 < <i>A</i> ≤ 35	56	
35 < <i>A</i> ≤ 45	34	
45 < <i>A</i> ≤ 55	19	
55 < <i>A</i> ≤ 65	7	

(a) Write down the modal class interval.


(1)

(b) Complete the cumulative frequency table.

(1)

(c) On the grid below, draw a cumulative frequency graph for your table.

(2)



(d) Use your graph to find an estimate for

(i) the median age of the employees,

.... years


(i) the interquartile range of the ages of the employees.

.....years

(3)

Another IT company has 80 employees. The age of the youngest employee is 24 years. The age of the oldest employee is 54 years. The median age is 38 years. The lower quartile age is 30 years. The upper quartile age is 44 years.

(e) On the grid below, draw a box plot to show information about the ages of the employees.



(Total 9 marks)

(2)

3. A company tested 100 batteries. The table shows information about the number of hours that the batteries lasted.


Time (t hours)	Frequency	Cumulative Frequency
50 ≤ <i>t</i> < 55	12	
55 ≤ <i>t</i> < 60	21	
60 ≤ <i>t</i> < 65	36	
65 ≤ <i>t</i> < 70	23	
70 ≤ <i>t</i> < 75	8	

(a) Complete the cumulative frequency table for this information.

(1)

(b) On the grid, draw a cumulative frequency graph for your completed table.

(2)



(c) Use your completed graph to find an estimate for the median time. You must state the units of your answer.

(2)

(Total 5 marks)


4. The table shows information about the ages of the 240 people at a club.

Age (t years)	Frequency	Cumulative Frequency
15 ≤ <i>t</i> < 20	95	
20 ≤ <i>t</i> < 25	90	
25 ≤ <i>t</i> < 30	35	
30 ≤ <i>t</i> < 35	15	
35 ≤ <i>t</i> < 40	5	

(a) Complete the cumulative frequency table.

(1)

(b) On the grid, draw the cumulative frequency graph for your table.



(c) Use your graph to find an estimate for the median age of the people.

...... years

(Total 4 marks)

(2)

5. An operator took 100 calls at a call centre. The table gives information about the time (*t* seconds) it took the operator to answer each call.


Time (t seconds)	Frequency	Cumulative Frequency
0 < <i>t</i> ≤ 10	16	
10 < <i>t</i> ≤ 20	34	
20 < <i>t</i> ≤ 30	32	
30 < <i>t</i> ≤ 40	14	
40 < <i>t</i> ≤ 50	4	

(a) Complete the cumulative frequency table.

(1)

(b) On the grid, draw a cumulative frequency graph for your table.

(2)




(c) Use your graph to find an estimate for the number of calls the operator took **more** than 18 seconds to answer.

(2)

(Total 5 marks)


6. 200 students took a test. The cumulative frequency graph gives information about their marks.



The lowest mark scored in the test was 10.

The highest mark scored in the test was 60.

Use this information and the cumulative frequency graph to draw a box plot showing information about the students' marks.



(Total 3 marks)


7. On Friday, Peter went to the airport.

He recorded the number of minutes that each plane was delayed.

He used his results to work out the information in this table.

	Minutes
Shortest delay	0
Lower quartile	2
Median	8
Upper quartile	18
Longest delay	41

(a) On the grid, draw a box plot to show the information in the table.




(2)

Peter also went to the airport on Saturday.

He recorded the number of minutes that each plane was delayed.

The box plot below was drawn using this information.



(b) Make two comparisons between the distributions of plane delays on Friday and on Saturday.

(Total 4 marks)

Histograms

Things to remember:

- Frequency = Frequency Density x Class Width;
- The y-axis will always be labelled "frequency density";
- The x-axis will have a continuous scale.


Questions:

1. One Monday, Victoria measured the time, in seconds, that individual birds spent on her bird table. She used this information to complete the frequency table.

Time (t seconds)	Frequency
0 < <i>t</i> ≤ 10	8
10 < <i>t</i> ≤ 20	16
20 < t ≤ 25	15
25 < <i>t</i> ≤ 30	12
30 < <i>t</i> ≤ 50	6


(a) Use the table to complete the histogram.





(3)

On Tuesday she conducted a similar survey and drew the following histogram from her results.




Time (Seconds)

(b) Use the histogram for Tuesday to complete the table.

Time (t seconds)	Frequency
0 < <i>t</i> ≤ 10	10
10 < <i>t</i> ≤ 20	
20 < <i>t</i> ≤ 25	
25 < <i>t</i> ≤ 30	
30 < <i>t</i> ≤ 50	

(2) (Total 5 marks)

This histogram gives information about the books sold in a bookshop one Saturday. 2.



(a) Use the histogram to complete the table.


Price (P) in pounds (£)	Frequency
0 < <i>P</i> ≤ 5	
5 < <i>P</i> ≤ 10	
10 < <i>P</i> ≤ 20	
20 < <i>P</i> ≤ 40	

(2)

The frequency table below gives information about the books sold in a second bookshop on the same Saturday.


Price (P) in pounds (£)	Frequency
0 < <i>P</i> ≤ 5	80
5 < <i>P</i> ≤ 10	20
10 < <i>P</i> ≤ 20	24
20 < <i>P</i> ≤ 40	96

(b) On the grid below, draw a histogram to represent the information about the books sold in the second bookshop.



(Total 5 marks)

3. The incomplete table and histogram give some information about the distances walked by some students in a school in one year.



(a) Use the information in the histogram to complete the frequency table.

Distance (d) in km	Frequency
0 < <i>d</i> ≤ 300	210
300 < <i>d</i> ≤ 400	350
400 < <i>d</i> ≤ 500	
500 < <i>d</i> ≤ 1000	

(2)

(b) Use the information in the table to complete the histogram.

(1)


(Total 3 marks)

4. The incomplete histogram and table show information about the weights of some containers.

Weight (w) in kg	Frequenc y
0 < <i>w</i> ≤ 1000	16
1000 < <i>w</i> ≤ 2000	
2000 < <i>w</i> ≤ 4000	
4000 < <i>w</i> ≤ 6000	16
6000 < <i>w</i> ≤ 8000	
8000 < <i>w</i> ≤ 12000	8

(a) Use the information in the histogram to complete the table.


(b) Use the information in the table to complete the histogram.



(2) (Total 4 marks)

(2)

5. The incomplete histogram and table give some information about the distances some teachers travel to school.



(a) Use the information in the histogram to complete the frequency table.

Distance (dkm)	Frequency
0 < <i>d</i> ≤ 5	15
5 < <i>d</i> ≤ 10	20
10 < <i>d</i> ≤ 20	
20 < <i>d</i> ≤ 40	
40 < <i>d</i> ≤ 60	10

(b) Use the information in the table to complete the histogram.


(1) (Total 3 marks)

(2)


6. The table gives information about the heights, in centimetres, of some 15 year old students.

Height (<i>h</i> cm) 145 < <i>h</i> ≤ 155		155 < <i>h</i> ≤ 175	175 < <i>h</i> ≤ 190	
Frequency	10	80	24	

Use the table to draw a histogram.



7. A teacher asked some year 10 students how long they spent doing homework each night. The histogram was drawn from this information.




Use the histogram to complete the table.

Time (t minutes)	Frequency
10 ≤ <i>t</i> < 15	10
15 ≤ <i>t</i> < 30	
30 ≤ <i>t</i> < 40	
$40 \le t < 50$	
50 ≤ <i>t</i> < 70	

(Total 2 marks)

Set Theory

Things to remember:




The intersection is where two sets overlap.

$$A \cap B$$

This means **A** and **B**.






If you put two sets together, you get the union.

$$A \cup B$$

This means **A** or **B**.





The **complement of A** is the region that is not A.

A'

This means **not A**.

Questions:

1.

$$\mathcal{E}$$
= {1, 2, 3, 4, 5, 6, 7, 8, 9, 10}

$$A = \{\text{multiples of 2}\}\$$

$$A \cap B = \{2, 6\}$$

$$A \cup B = \{1, 2, 3, 4, 6, 8, 9, 10\}$$

Draw a Venn diagram for this information.


(Total for question is 4 marks)

- **2.** Here is a Venn diagram.
 - (a) Write down the numbers that are in set
 - (i) *A* ∪ *B*

.....

(ii) $A \cap B$

(2)



One of the numbers in the diagram is chosen at random.

(b) Find the probability that the number is in set A'

(2)

(Total for question = 4 marks)

All 5 19 p 16 p 21 p 24 p 40 p	ni asked 50 people which drinks they liked from tea, coffee and milk. 0 people like at least one of the drinks eople like all three drinks. eople like tea and coffee but do not like milk. eople like coffee and milk. eople like tea and milk. eople like tea and milk. eople like coffee. erson likes only milk.
Sam (a)	ni selects at random one of the 50 people. Work out the probability that this person likes tea.
(b)	Given that the person selected at random from the 50 people likes tea, find the probability that this person also likes exactly one other drink.
	(2) (Total for question = 6 marks)

3.

Proportion

Things to remember:

- Start by checking the question for squares, cubes and roots;
- "x is directly proportional to y" looks like x α y or x = ky
- "x is inversely proportional to y" looks like $\mathbf{x} \propto \frac{1}{y}$ or $\mathbf{x} = \frac{k}{y}$

Q		Δ	c	ŧi	^	n	c	
u	u	u	5	LI	u	H	2	

1.	The shutter speed, S, of a camera varies inversely as the square of the aperture setting, f.
	When $f = 8$, $S = 125$

((a)	Find a	a formula	for	Sin	terms	of	f.

 ٠.			 													 									
																						(:	3)

(b) Hence, or otherwise, calculate the value of S when f = 4

(Total 4 marks)

2. In a factory, chemical reactions are carried out in spherical containers. The time, *T* minutes, the chemical reaction takes is directly proportional to the square of the radius, *R* cm, of the spherical container.

When R = 120, T = 32Find the value of T when R = 150

	irectly proportional to the square of t . 0 when $t = 4$	
	Express <i>d</i> in terms of <i>t</i> .	(a)
(3)	Work out the value of d when $t = 7$	(b)
(1)	d = Work out the positive value of t when $d = 45$	(c)
(2) (Total 6 marks)	$t = \dots$	
tional to the square of the time, t,	distance, D , travelled by a particle is directly proport t . When $t = 40$, $D = 30$ Find a formula for D in terms of t .	
(3)	D = Calculate the value of D when $t = 64$	(b)
(1)	Calculate the value of t when $D = 12$ Give your answer correct to 3 significant figures.	(c)
(2) (Total 6 marks)		

• • •	The time, T seconds, it takes a water heater to boil so to the mass of water, m kg, in the water heater. Wher (a) Find T when $m = 400$	5.
$T = \dots $ (3) a constant mass of water is inversely	T: The time, T seconds, it takes a water heater to boil a	
	proportional to the power, P watts, of the water heate When $P = 1400$, $T = 360$ (b) Find the value of T when $P = 900$	
「=(3) (Total 6 marks)	T =	
	A ball falls vertically after being dropped. The ball falls a distance <i>d</i> metres in a time of <i>t</i> second is directly proportional to the square of <i>t</i> . The ball falls 20 metres in a time of 2 seconds. (a) Find a formula for <i>d</i> in terms of <i>t</i> .	6.
d =(3)	denotes the distance the ball falls in 3 second	
m (1) n.	(c) Calculate the time the ball takes to fall 605 m.	
seconds (3) (Total 7 marks)		

າ (x cm). When the	a spring, the tension (<i>T</i> newtons) is directly proportional to its extension sion is 150 newtons, the extension is 6 cm. Find a formula for <i>T</i> in terms of <i>x</i> .	
(3)	T =	(t
newtons (1)	Calculate the extension, in cm, when the tension is 600 newtons.	(0
cm (1) (Total 5 marks)	inversely proportional to d . en $d = 50$, $f = 256$ d the value of f when $d = 80$	V
(Total 3 marks)	f =	

Percentages – compound interest


Thing		emember:		
•	new a	$amount = original amount x multiplier^n $	Number of years	
Ques 1.	At the	y invests £4500 at a compound interest rate of see end of <i>n</i> complete years the investment has gothe value of <i>n</i> .	5% per annum.	
				(Total 2 marks)
2.		uys a new machine. value of the machine depreciates by 20% each y Bill says 'after 5 years the machine will have r Bill is wrong. Explain why.	no value'.	
	Bill w	ants to work out the value of the machine after By what single decimal number should Bill munew?		(1)
				(2) (Total 3 marks)
3.		n bought a new car. Each year, the value of her ulate the number of years after which the value		
4.	The v	value of a car depreciates by 35% each year.		(Total 3 marks)
T.	At the	e end of 2007 the value of the car was £5460 out the value of the car at the end of 2006		
			£	
				(Total 3 marks)

5.	Toby invested £4500 for 2 years in a savings account. He was paid 4% per annum compound interest. (a) How much did Toby have in his savings account after 2 years?
	\pounds
	(b) Work out the value of n. (7) (Total 5 marks
6.	Mario invests £2000 for 3 years at 5% per annum compound interest. Calculate the value of the investment at the end of 3 years.
	£(Total 3 marks

Percentages - reverse

Things to remember:

• Work out what the multiplier would have been;



_							
Q	Δ	c.	tı.	\smallfrown	n	c	
w	•			u			_

1.	Loft insulation reduces annual heating costs by 20%.
	After he insulated his loft, Curtley's annual heating cost was £520.
	Work out Curtley's annual heating cost would have been, if he had not insulated his loft.

£	
	(Total 3 marks)

2. In a sale, normal prices are reduced by 20%.

SALE - 20% OFF

Andrew bought a saddle for his horse in the sale.

The sale price of the saddle was £220.

Calculate the normal price of the saddle.

£													 										 											
														((1	Γ	C)	t	а	l	3	}	r	n	1	a	r	k	(S)	

3. Hajra's weekly pay this year is £240
This is 20% more than her weekly pay last year.
Bill says 'This means Hajra's weekly pay last year was £192'.
Bill is wrong,

(a) Explain why.

(1)

(b) Work out Hajra's weekly pay last year.

£(2)

(Total 3 marks)

4.	The p (a)	rice of all rail season tickets to London increas The price of a rail season ticket from Cambrid Work out the price before this increase.	
			£(2
	(b)	After the increase, the price of a rail season to £2828.80 Work out the price before this increase.	
			£
			(3) (Total 5 marks)
5.	The s	ale, normal prices are reduced by 25%. ale price of a saw is £12.75 late the normal price of the saw.	
			£
6.	The s	ale, normal prices are reduced by 12%. ale price of a DVD player is £242. out the normal price of the DVD player.	(Total 3 marks)
			£(Total 3 marks
7.	It offe Dave	age sells cars. rs a discount of 20% off the normal price for ca pays £5200 cash for a car. late the normal price of the car.	ash.
			£(Total 3 marks

Useful websites:

www.mathswatchvle.com

www.methodmaths.com

www.hegartymaths.com

www.mymaths.co.uk

www.drfrost.com

www.bbc.co.uk/schools/gcsebitesize /maths

Remember: Do your best; it is all you can do ©